

ANALYTICAL SOLUTIONS OF SOME SPECIAL NONLINEAR PARTIAL DIFFERENTIAL EQUATIONS USING ELZAKI-ADOMIAN DECOMPOSITION METHOD

¹Orapine Hycienth O., ^{1*}Baidu Ali A., ²Oko Godfrey, ¹Nyamtwam Elija Vafa

¹Department of Mathematics, Nigerian Army University Biu, Borno State, Nigeria

²Department of Mathematical Sciences, Nigerian Defence Academy, Kaduna

*Corresponding Author Email Address: aliaudubaidu@yahoo.com

ABSTRACT

We apply the Elzaki-Adomian Decomposition Method (EADM) in this study to solve nonlinear Benjamin-Bona-Mahony (BBM) and Fisher's partial differential equations (PDE). This method, being an integral transform, is a hybrid of two well-known and efficient methods: the Elzaki transform and the Adomian decomposition method. The method is demonstrated by solving two special cases of the BBM Equation and one special case of Fisher's partial differential equation. Because of its high convergence rate in approximating exact solutions, this approach is very dependable. The method can also produce numerical solutions without the usage of restrictive assumptions or the discretization typical of numerical methods; making it free of round-off errors. The Elzaki-Adomian Decomposition method employs a straightforward computation that leads to effectiveness. The efficiency of EADM is demonstrated in the significant reduction of number of numerical computations. The effectiveness and efficiency of EADM account for its broad application, particularly for higher order PDEs.

Keywords: Elzaki Transform, Adomian Polynomials, PDEs, Analytical Solutions, Benjamin-Bona-Mahony Equation, Fisher's Equation

INTRODUCTION

Nonlinear differential equations are incredibly essential to humans since most physical phenomena are nonlinear in nature and are modelled by these equations. In this regard, partial differential equations (PDEs) are in particular, fundamental. Unfortunately, analytical techniques cannot be used to solve the majority of nonlinear problems. In addition, to solve nonlinear problems, standard numerical methods require perturbation, discretization, linearization, or transformation. However, Adomian (1994) established that the Adomian decomposition method is free of such steps as are involved in standard numerical methods and is thus widely used in the literature. Several researchers have made significant efforts and implemented diverse methods for solving nonlinear PDEs over the last few decades. Recently Ali *et al.*, (2018) applied the Laplace Adomian decomposition method in finding the approximate solutions for nonlinear general fisher's equation. In a similar vein, we reference the works of (Khuri, 2001; Wazwaz, 2010; Wazwaz and Mehanna, 2010).

Another important method that has received little attention is the Elzaki-Adomian decomposition method (EADM). It was introduced by Ige, *et al.*, (2019). It is a combination of the Elzaki transform and the Adomian decomposition method, two well-known and efficient methods. It is possible to obtain numerical solutions using this method without the use of restrictive assumptions or discretization,

making it free of round-off errors. A solution in the form of a finite series is also achieved using this method, and it has the highest and fastest rate of convergence. In this paper we apply EADM to obtain the analytical solution of some special nonlinear partial differential equations: The Benjamin-Bona-Mahony (BBM) Equations and Fisher's Equations.

The Benjamin-Bona-Mahony (BBM) Equations

The partial differential equation Benjamin-Bona-Mahony (BBM), commonly known as the regularized long-Wave equation (RLWE), was introduced by Benjamin *et al.* (1972). (See also Muhammad *et al.*, 2019).

$$U_t + U_x + UU_x - U_{xxt} = 0, \quad U(x, 0) = f(x) \quad (1.0)$$

Benjamin, Bona, and Mahony explored this equation in 1972 as an improvement on the Korteweg de Vries equation (KdV equation) for modeling long surface gravity waves of small amplitude traveling in 1+1 dimensions. They demonstrated the BBM equation's solutions' stability and uniqueness. The KdV equation, on the other hand, is unstable in its high wavenumber components. Furthermore, the KdV equation has an infinite number of motion integrals, whereas the BBM equation has just three (Molati and Khalique, 2012)

In physical applications, the BBM equation is well-known. It offers a model for long-wave propagation that includes nonlinear and dissipative phenomena. It's used to study long-wavelength surface waves in liquids, cold plasma hydro magnetic waves, compressible fluids acoustic-gravity waves, and harmonic crystal acoustic waves (Molati and Khalique, 2012). The dynamics of the BBM equation has drawn the attention of many mathematicians (Singh *et al.*, 2011). For shallow water waves, the BBM equation has been examined as a regularized version of the Kdv equation. Finding analytic solutions to the nonlinear BBM Equation is crucial, as the equation also models complicated physical systems that can occur in engineering, chemistry, biology, mechanics, and physics (Talha and Khaled, 2009).

Fisher's Equations

As a nonlinear model for a physical system comprising linear diffusion and nonlinear growth, the Fisher equation assumes the following non-dimensional form:

$$U_t = U_{xx} + \mu(1 - U^\alpha)(U - \rho), \quad U(x, 0) = g(x) \quad (1.2)$$

A constant-velocity front of transition from one homogeneous

condition to another is described by (1.2) kink-like traveling wave solutions called Solitons. Solitons, on the other hand, emerge as a result of a delicate balancing between weak nonlinearity and dispersion. As a result, in Mathematics and Physics, a soliton is defined as a self-reinforcing solitary wave—a wave packet or pulse that keeps its shape while traveling at steady velocity. The dispersion relation between the frequency and the speed of the waves is referred to as "dispersive effects." Solitons are solutions to a class of weakly nonlinear dispersive partial differential equations that describe physical systems. Instead of dispersion, when diffusion occurs, energy released by nonlinearity balances energy consumed by diffusion, resulting in moving waves or fronts. As a consequence, moving wave fronts are a well-studied solution form for reaction diffusion equations, with applications in chemistry, biology, and medicine (Wazwaz and Gorguis, 2004).

Description of Elzaki Transform

Elzaki transform is an integral transformation defined for function of exponential order (Tarig, 2011a). Consider the function in the set A defined as;

$$A = \left\{ f(t): \exists M, c_1, c_2 > 0, |f(t)| < M e^{\frac{|t|}{c_1}}, \text{ if } t \in (-1)^i \times [0, \infty) \right\} \quad (2.1)$$

where for any given function in the set A defined above, the constant c_1, c_2 may be either finite or infinite, but M must be infinite. According to Tarig (2011a), Elzaki Transform is defined as:

$$E\{f(t)\} = w^2 \int_0^\infty f(wt)e^{-t} dt = T(w), \quad t \geq 0, \quad w \in (c_1, c_2) \quad (2.2)$$

Or

$$E\{f(t)\} = w \int_0^\infty f(t)e^{-\frac{t}{w}} dt = T(w), \quad t \geq 0, \quad w \in (c_1, c_2) \quad (2.3)$$

We note here, that w in the above definition is used to factor t in the analysis of function f .

Elzaki Transform of Partial Derivatives Tarig *et al.* (2011b) extended the method to solving partial differential equations. The Elzaki transform of partial derivatives are obtained through integration by parts, then we find the following expressions;

$$E \left[\frac{\partial f(x, t)}{\partial t} \right] = \frac{T(x, w)}{w} - wf(x, 0) \quad (2.4)$$

$$E \left[\frac{\partial^2 f(x, t)}{\partial t^2} \right] = \frac{T(x, w)}{w^2} - f(x, 0) - w \frac{\partial f(x, 0)}{\partial t} \quad (2.5)$$

$$E \left[\frac{\partial^3 f(x, t)}{\partial t^3} \right] = \frac{T(x, w)}{w^3} - \frac{f(x, 0)}{w} - \frac{\partial f(x, 0)}{\partial t} \quad (2.6)$$

$$E \left[\frac{\partial f(x, t)}{\partial t} \right] = \frac{d}{dx} [T(x, w)] \quad (2.7)$$

$$E \left[\frac{\partial^2 f(x, t)}{\partial x^2} \right] = \frac{d^2}{dx^2} [T(x, w)] \quad (2.8)$$

$$E \left[\frac{\partial^3 f(x, t)}{\partial x^3} \right] = \frac{d^3}{dx^3} [T(x, w)] \quad (2.9)$$

Elzaki Transform of Some Functions

By using the definition of Elzaki transform of equations (2.2)-(2.3) on some functions the results can be generated as tabulated in

table 1. (Tarig, 2011a).

Table 1: Table of Functions and their Elzaki Transform

$f(t)$	$E[f(t)] = T(w)$
1	w^2
t	w^3
t^n	$n! w^{n+2}$
$\frac{t^{a-1}}{\Gamma(a)}, \quad a > 0$	w^{a+1}
e^{at}	$\frac{w^2}{1-aw}$
$t e^{at}$	$\frac{w^2}{w^3}$
$\frac{t^{n-1} e^{at}}{(n-1)!}, n = 1, 2, \dots$	$\frac{(1-aw)^2}{w^{n+1}}$
$\sin at$	$\frac{(1-aw)^n}{(1-aw)^n}$
$\cos at$	$\frac{aw^3}{aw^3}$
$\sinh at$	$\frac{1+a^2w^2}{w^2}$
$\cosh at$	$\frac{1+a^2w^2}{aw^2}$
$e^{at} \sin bt$	$\frac{1-a^2w^2}{1-a^2w^2}$
$e^{at} \cos bt$	$\frac{bu^3}{(1-aw)^2 + b^2w^2}$
$t \sin at$	$\frac{(1-aw)w^2}{(1-aw)^2 + b^2w^2}$
$J_0(at)$	$\frac{2aw^2}{2aw^2}$
$H(t-a)$	$\frac{1+a^2w^2}{w^2}$
$\delta(t-a)$	$\frac{\sqrt{1+a^2w^2}}{w^2 e^{\frac{a}{w}}}$
	$w e^{\frac{a}{w}}$

METHODOLOGY

In this paper, our interest is to solve some special nonlinear partial differential equations which are third order Benjamin-Bona-Mahony and second order Fisher's equations. We first demonstrate how the Elzaki transform method can be used to decompose the general nonlinear partial differential equation.

According to Ziane and Hamdi (2015), we consider;

$$\frac{\partial^n u(x, t)}{\partial t^n} + Ru(x, t) + Nu(x, t) = g(x, t) \quad (3.1)$$

where $n = 1, 2, 3$.

And the initial condition is given as

$$\frac{\partial^{n-1} u(x, t)}{\partial t^{n-1}} \Big|_{t=0} = f_{n-1}(x)$$

where $\frac{\partial^n u(x, t)}{\partial t^n}$ is the partial derivative of function $u(x, t)$ of n th order, while R represents the linear differential operator, $Nu(x, t)$ represents the nonlinear terms of the differential equations, and $f(x, t)$ indicates the non-homogeneous (source) term.

Applying the Elzaki transform on equation (3.1) we have;

$$E \left[\frac{\partial^n u(x, t)}{\partial t^n} \right] + E[Ru(x, t)] = E[g(x, t)] \quad (3.2)$$

We recall that

$$E \left[\frac{\partial^n u(x, t)}{\partial t^n} \right] = \frac{E[u(x, t)]}{w^n} - \sum_{k=0}^{n-1} w^{2-n+k} \frac{\partial^k u(x, 0)}{\partial t^k} \quad (3.3)$$

substituting Equation (3.3) into Equation (3.2), we have;

$$E[u(x, t)] = w^n E[g(x, t)] + \sum_{k=0}^{n-1} w^{2+k} \frac{\partial^k u(x, 0)}{\partial t^k} - w^n \{E[Ru(x, t)] + E[Nu(x, t)]\} \quad (3.3)$$

Applying the inverse Elzaki transform to Equation (3.3), we have;

$$u(x, t) = E^{-1} \left[w^n E[g(x, t)] + \sum_{k=0}^{n-1} w^{2+k} \frac{\partial^k u(x, 0)}{\partial t^k} - E^{-1} \{w^n \{E[Ru(x, t)] + E[Nu(x, t)]\} \right] \quad (3.4)$$

We can rewrite this as;

$$u(x, t) = F(x, t) - E^{-1} \{w^n \{E[Ru(x, t)] + E[Nu(x, t)]\} \} \quad (3.5)$$

where $F(x, t)$ represents the expression that rises from the given initial conditions and the source terms after simplification. We note here that, our solution will be in the form of infinite series as

$$u(x, t) = \sum_{k=0}^{\infty} u_k(x, t) \quad (3.6)$$

We can now decompose the nonlinear term as

$$Nu(x, t) = \sum_{k=0}^{\infty} A_k \quad (3.7)$$

Where A_k is defined as the Adomian polynomials which can be generated using the formula

$$A_k = \frac{1}{k!} \frac{\partial^k}{\partial y^k} \left[N \left(\sum_{j=0}^{\infty} \rho^j u_j \right) \right]_{\rho=0}, k = 0, 1, \dots, (3.8a)$$

where ρ is taken as formal parameter and we will drop it after the calculation by equating it to zero.

According to El-Kalla (2007), Adomian polynomial can be computed in different ways it is not unique and we can calculate it from the Tylor expansion of function $f(u)$ around the first component u_0 i.e.

$$f(u) = \sum_{k=0}^{\infty} A_k = \sum_{k=0}^{\infty} \frac{(u - u_0)^k}{k!} f^{(k)}(u_0) \quad (3.8b)$$

Here, we provided the first five Adomian polynomials for the nonlinear terms $Nu = f(u)$

$$A_0 = f(u_0)$$

$$A_1 = u_1 f'(u_0),$$

$$A_2 = u_2 f'(u_0) + \frac{u_1^2}{2!} f''(u_0),$$

$$A_3 = u_3 f'(u_0) + u_1 u_2 f''(u_0) + \frac{u_1^3}{3!} f'''(u_0),$$

$$A_4 = u_4 f'(u_0) + \left(u_1 u_3 + \frac{u_2^2}{2!} \right) f''(u_0) + \left(\frac{u_1^2 u_2}{2!} \right) f'''(u_0) + \frac{u_1^4}{4!} f^{(4)}(u_0),$$

Now substituting Equation (3.7) and Equation (3.6) into Equation (3.5) we have that;

$$\sum_{k=0}^{\infty} u_k(x, t) = F(x, t) - E^{-1} \left\{ w^n \left\{ E \left[R \sum_{k=0}^{\infty} u_k(x, t) \right] + E \left[\sum_{k=0}^{\infty} A_k \right] \right\} \right\} \quad (3.9)$$

Then starting to evaluate from equation (3.9) at $k = 0$, we have

$$u_0(x, t) = F(x, t) \quad (3.10)$$

And the recursive relation from equation (3.9) given as:

$$u_{k+1} = -E^{-1} \{w^n \{E[Ru_k(x, t)] + E[A_k]\} \} \quad (3.11)$$

where $n = 1, 2, 3$ (from the order of the PDE) and $k \geq 0$. The analytical solution $u(x, t)$ can be approximated by truncated series.

$$u(x, t) = \lim_{k \rightarrow \infty} \sum_{k=0}^{\infty} u_k(x, t) \quad (3.12)$$

The infinite series in equation (3.12) may converge completely very fast to exact solution or with few terms truncation will result to the exact solution of the given differential equation.

Numerical Implementations

Numerical Problem 1

We consider the BBM equation,

$$U_t + U_x + UU_x - U_{xxt} = 0, \quad U(x, 0) = x \quad (4.1)$$

Taking Elzaki transform of each term we have;

$$E[U_t] = E[U_{xxt}] - E[U_x + UU_x] \quad (4.2)$$

where

$$E[U_t] = \frac{E[U(x, t)]}{w} - wU(x, 0)$$

Thus Equation (4.2) becomes;

$$E[U(x, t)] = w^2 U(x, 0) + w \cdot E[U_{xxt} - U_x - UU_x] \quad (4.3)$$

Introducing the initial condition and taking the inverse of the Elzaki transform we have;

$$U(x, t) = x + E^{-1} \{w \cdot E[U_{xxt} - U_x - UU_x]\} \quad (4.4)$$

But

$$U(x, t) = \sum_{k=0}^{\infty} U_k(x, t) \quad (4.5)$$

and

$$UU_x = \sum_{k=0}^{\infty} A_k \quad (4.6)$$

Substituting Equations (4.5)-(4.6) in Equation (4.4) we have

$$U_0 = x \quad (4.7)$$

and the recurrence relation as

$$U_{k+1} = E^{-1} \{w \cdot E[U_{kxxt} - U_{kx} - A_k]\} \quad (4.8)$$

Now we compute the individual terms from the recurrence equation.

$$U_1 = E^{-1}[w \cdot E [U_{0xxt} - U_{0x} - A_0]] \quad (4.9)$$

where,

$$A_0 = U_0 U_{0x} = x \cdot 1 = x, \quad U_{0xxt} = 0 \quad \text{and} \quad U_{0x} = 1$$

Thus Equation (4.9) becomes

$$\begin{aligned} U_1 &= E^{-1}[w \cdot E [0 - 1 - x]] \\ &= E^{-1}[w^3 \cdot (-1 - x)] \\ U_1 &= -(1 + x)t \end{aligned} \quad (4.10)$$

Again from equation (4.8) we have;

$$U_2 = E^{-1}[w \cdot E [U_{1xxt} - U_{1x} - A_1]] \quad (4.11)$$

where,

$$\begin{aligned} A_1 &= U_0 \frac{\partial U_1}{\partial x} + U_1 \frac{\partial U_0}{\partial x} \\ &= x(-t) - (1 + x)t \cdot 1 \\ &= -(2x + 1)t \end{aligned}$$

Thus, it follows from equation (4.11) that;

$$U_2 = (x + 1)t^2 \quad (4.12)$$

Similarly, from equation (4.8) we compute U_3 as follows;

$$U_3 = E^{-1}[w \cdot E [U_{2xxt} - U_{2x} - A_2]] \quad (4.13)$$

Where;

$$\begin{aligned} A_2 &= U_0 \frac{\partial U_2}{\partial x} + U_1 \frac{\partial U_1}{\partial x} + U_2 \frac{\partial U_0}{\partial x} \\ &= (3x + 2)t^2 \end{aligned}$$

Hence equation (4.13) becomes;

$$\begin{aligned} U_3 &= \frac{2!}{3!} \cdot (-3(x + 1)) \cdot E^{-1}[3! w^5] \\ U_3 &= -(x + 1)t^3 \end{aligned} \quad (4.14)$$

Therefore, the solution of equation (4.1) is

$$U(x, t) = U_0 + U_1 + U_2 + U_3 + \dots \quad (4.15)$$

$$\begin{aligned} &= \frac{x}{1} + \left[\frac{-(1 + x)t}{(1 + t)} \right] \\ &= \frac{x(1 + t) - (1 + x)t}{1 + t} \end{aligned}$$

$$U(x, t) = \frac{x - t}{1 + t} \quad (4.16)$$

Clearly, Equation (4.16) can be shown to be the exact solution of the BBM Equation (4.1) and its plot shown in fig.1.




Fig.1: Plot of the solution of BBM equation (4.1) by Elzaki-Adomian Decomposition Method.

Numerical Problem 2

We consider the BBM equation (Shehata, 2015).

$$U_t = U_{xxt} - U_x - UU_x \quad (4.17)$$

subject to the initial condition

$$U(x, 0) = \operatorname{sech}^2\left(\frac{x}{4}\right) \quad (4.18)$$

with the exact solution

$$U(x, t) = \operatorname{sech}^2\left(\frac{x}{4} - \frac{t}{3}\right) \quad (4.19)$$

Similarly, taking Elzaki transform of each term we have;

$$E[U_t] = E[U_{xxt}] - E[U_x + UU_x] \quad (4.20)$$

Using equation (2.3), thus Equation (4.20) becomes;

$$E[U(x, t)] = w^2 U(x, 0) + w \cdot E[U_{xxt} - U_x - UU_x] \quad (4.21)$$

Applying the initial condition and taking the inverse transform we have;

$$\begin{aligned} U(x, t) &= \operatorname{sech}^2\left(\frac{x}{4}\right) \\ &+ E^{-1}[w \\ &\cdot E[U_{xxt} - U_x - UU_x]] \end{aligned} \quad (4.22)$$

But

$$U(x, t) = \sum_{k=0}^{\infty} U_k(x, t) \quad (4.23)$$

and

$$UU_x = \sum_{k=0}^{\infty} A_k \quad (4.24)$$

Putting Equation (4.23) and (4.24) in Equation (4.22) we have;

$$U_0 = \operatorname{sech}^2\left(\frac{x}{4}\right) \quad (4.25)$$

and the recurrence equation as

$$U_{k+1} = E^{-1}[w \cdot E [U_{kxxt} - U_{kx} - A_k]] \quad (4.26)$$

Similarly, we calculate the individual terms from the series solution from the equation (4.26)

$$U_1 = E^{-1}[w \cdot E [U_{0xxt} - U_{0x} - A_0]] \quad (4.27)$$

Where,

$$U_{0x} = -\frac{\sinh\left(\frac{x}{4}\right)}{2 \cosh^3\left(\frac{x}{4}\right)}, \quad U_{0xxt} = 0 \quad (4.28)$$

$$A_0 = U_0 U_{0x} = -\frac{\sinh\left(\frac{x}{4}\right)}{2 \cosh^5\left(\frac{x}{4}\right)} \quad (4.29)$$

Thus Equation (4.27) becomes

$$U_1 = \frac{t}{2} \sinh\left(\frac{x}{4}\right) \left[\operatorname{sech}^3\left(\frac{x}{4}\right) + \operatorname{sech}^5\left(\frac{x}{4}\right) \right] \quad (4.30)$$

Again from equation (4.26) we have;

$$U_2 = E^{-1}[w \cdot E [U_{1xxt}] - w \cdot E [U_{1x} + A_1]] \quad (4.31)$$

$$U_{1x} = \frac{(2 \cosh^4\left(\frac{x}{4}\right) + \cosh^2\left(\frac{x}{4}\right) - 5)t}{8 \cosh^6\left(\frac{x}{4}\right)} \quad (4.32)$$

$$\begin{aligned} &U_{1xxt} \\ &= \frac{(2 \cosh^4\left(\frac{x}{4}\right) + 2 \cosh^2\left(\frac{x}{4}\right) - 15) \sinh\left(\frac{x}{4}\right)}{16 \cosh^7\left(\frac{x}{4}\right)} \end{aligned} \quad (4.33)$$

$$A_1 = -\frac{(4 \cosh^4\left(\frac{x}{4}\right) + \cosh^2\left(\frac{x}{4}\right) - 7)t}{8 \cosh^8\left(\frac{x}{4}\right)} \quad (4.34)$$

Plugging equations (4.32)-(4.34) into equations (4.31) and simplifying we obtain;

$$U_2 = \frac{t \left(2\cosh^4\left(\frac{x}{4}\right) + 2\cosh^2\left(\frac{x}{4}\right) - 15 \right) \sinh\left(\frac{x}{4}\right)}{16\cosh^7\left(\frac{x}{4}\right)} + \frac{\left(2\cosh^6\left(\frac{x}{4}\right) + 5\cosh^4\left(\frac{x}{4}\right) - 4\cosh^2\left(\frac{x}{4}\right) - 7 \right) t^2}{8\cosh^8\left(\frac{x}{4}\right)}$$

And using trigonometry identities we have

$$U_2 = \frac{t}{16} \sinh\left(\frac{x}{4}\right) \left(2\operatorname{sech}^3\left(\frac{x}{4}\right) + 2\operatorname{sech}^5\left(\frac{x}{4}\right) - 15 \operatorname{sech}^7\left(\frac{x}{4}\right) \right) + \left(4\operatorname{sech}^2\left(\frac{x}{4}\right) + \frac{5}{8} \operatorname{sech}^4\left(\frac{x}{4}\right) - 2\operatorname{sech}^6\left(\frac{x}{4}\right) - \frac{7}{8} \operatorname{sech}^8\left(\frac{x}{4}\right) \right) t^2 \quad (4.36)$$

Hence the approximate solution of the BBM equations (4.17-4.18) is

$$U(x, t) = U_0 + U_1 + U_2 + \dots \quad (4.37)$$

$$U(x, t) = \operatorname{sech}^2\left(\frac{x}{4}\right) + \frac{t}{2} \sinh\left(\frac{x}{4}\right) \left[\operatorname{sech}^3\left(\frac{x}{4}\right) + \operatorname{sech}^5\left(\frac{x}{4}\right) \right] + \frac{t}{16} \sinh\left(\frac{x}{4}\right) \left(2\operatorname{sech}^3\left(\frac{x}{4}\right) + 2\operatorname{sech}^5\left(\frac{x}{4}\right) - 15 \operatorname{sech}^7\left(\frac{x}{4}\right) \right) + \left(4\operatorname{sech}^2\left(\frac{x}{4}\right) + \frac{5}{8} \operatorname{sech}^4\left(\frac{x}{4}\right) - 2\operatorname{sech}^6\left(\frac{x}{4}\right) - \frac{7}{8} \operatorname{sech}^8\left(\frac{x}{4}\right) \right) t^2 + \dots \quad (4.38)$$




Fig.2: Plot of the exact solution of BBM equations (4.17)-(4.18)




Fig. 3: Plot of the solution of BBM equations (4.17)-(4.18) by Elzaki-Adomian Decomposition Method




Fig. 4: Plot of the solution of BBM equations (4.17)-(4.18) by Adomian Decomposition Method (Shehata, 2015)

Numerical Problem 3

Consider the fisher Equation (Ali *et al.*, 2018).

$$U_t = U_{xx} + U - U^2, \quad U(x, 0) = \beta \quad (4.39)$$

Taking Elzaki transform on each term we have;

$$E[U(x, t)] = w^2 U(x, 0) + w \cdot E[U_{xx} + U - U^2] \quad (4.40)$$

Applying the initial condition and the taking the inverse transform we have;

$$U(x, t) = \beta + E^{-1}[w \cdot E[U_{xx} + U - U^2]] \quad (4.41)$$

But

$$U(x, t) = \sum_{n=0}^{\infty} U_n(x, t)$$

$$U^2 = \sum_{n=0}^{\infty} A_n$$

Thus, from equation (4.41) we obtain

$$\sum_{n=0}^{\infty} U_n(x, t) = \beta + E^{-1} \left[w \cdot E \left[\sum_{n=0}^{\infty} U_{nxx} + \sum_{n=0}^{\infty} U_n - \sum_{n=0}^{\infty} A_n \right] \right] \quad (4.41)$$

where

$$U_0 = \beta \quad (4.42)$$

And the recurrence relation is given as

$$U_{n+1} = E^{-1} [w \cdot E [U_{nxx} + U_n - A_n]] \quad (4.43)$$

Thus, the remaining terms of the series solution for the equation (4.39) are calculated.

$$U_1 = E^{-1} [w \cdot E [U_{0xx} + U_0 - A_0]] \quad (4.44)$$

where $A_0 = U_0^2 = \beta^2$ and thus;

$$U_1 = (\beta - \beta^2)t \quad (4.45)$$

In the same vein, from equation (4.43) we compute

$$U_2 = E^{-1} [w \cdot E [U_{1xx} + U_1 - A_1]] \quad (4.46)$$

where

$$A_1 = 2U_1U_0 = 2(\beta^2 - \beta^3)t$$

and thus;

$$U_2 = \frac{t^2}{2!} \beta(1 - \beta)(1 - 2\beta) \quad (4.47)$$

Thus, it follows that;

$$U(x, t) = U_0 + U_1 + U_2 + \dots \quad (4.48)$$

and hence;

$$U(x, t) = \beta + \beta(1 - \beta)t + \beta(1 - \beta)(1 - 2\beta) \frac{t^2}{2!} + \dots \quad (4.49)$$

After some algebraic manipulations and simplification, we have the exact solution to the equation (4.39)

$$U(x, t) = \frac{\beta e^t}{1 - \beta + \beta e^t} \quad (4.50)$$




Fig. 5: Plot of the solution of fisher's equation (4.39) using Elzaki-Adomian Decomposition Method.

Conclusion

In this paper, the Elzaki-Adomian Decomposition Method (EADM) has been successfully applied to find the solutions of nonlinear Benjamin-Bona-Mahony and Fisher's equations as presented in figures 1-5. It is observed that the use of hybrid EADM provides very good approximate solutions when compared with exact values

than Adomian Decomposition Method (ADM). The method transforms these equations to recurrences relation whose terms can be computed with the aid of any symbolic computational environment such as Maple, Mathematica, and Scientific workplace among others. The solution using this method is usually in the form of a finite series and it has high and fastest rate of convergence to the exact solutions of the relevant problems. It is possible to obtain numerical solutions using this method without the use of restrictive assumptions or discretization, making it free of round-off errors. The Elzaki-Adomian Decomposition method use a simple and straightforward calculation. The number of numerical computations is decreased. The efficiency of EADM and the reduction in calculations demonstrate its extensive applicability, particularly for higher order PDEs.

REFERENCES

- Adomian G. (1994). Solving frontier problems of physics: the decomposition method, *Kluwer Academic, Dordrecht*.
- Ali A, Humaira, Laila, et al., (2018). Analytical solution of General Fisher's Equation by using Laplace Adomian decomposition method. *J Pur Appl Math*. 2(3):01-4.
- Ali A. Humaira, Laila, and Kamal shah (2018) Analytical solution of General Fisher's Equation by using Laplace Adomian decomposition method. *Journal of Pure and Applied Mathematics*. 2(3): 01-4.
- Benjamin, T. B., Bona, J. L. and Mahony, J. J. (1972). Model equations for long waves in nonlinear dispersive systems, *Philos. Trans. Royal Soc., London A*, 272: 47-78
- Djelloul Ziane and Hamdi Cherif M. (2015). Resolution of nonlinear partial differential equations by elzaki transform decomposition method. *Journal of Approximation Theory and Applied Mathematics* 5:17-30.
- El-Kalla I. L. (2007). Error analysis of adomian series solution to a class of nonlinear differential equations. *Applied Mathematics E-Notes*, 7:214-221.
- Hendi FA. (2011). Laplace Adomian decomposition method for solving the nonlinear volterra integral equation with weakly kernels, *Studies in Nonlinear Sciences*. 2:129-34.
- Ige, O., Oderinu, R., Elzaki, T. (2019). 'Adomian Polynomial and Elzaki Transform Method of Solving Fifth Order Korteweg-De Vries Equation', *Caspian Journal of Mathematical Sciences (CJMS)*, 8(2):103-119. doi:10.22080/cjms.2018.14486.1346
- Khuri SA. (2001). A Laplace decomposition algorithm applied to a class of nonlinear differential equations. *J Math Annl Appl*. 1(4):141-55.
- Molati, M., and Khalique, C.M. (2012). Lie symmetry analysis of the time-variable coefficient BBM equation. *Advances in Difference Equations*, 212. doi:10.1186/1687-1847-2012-212.
- Muhammad Ikram, Abbas Muhammad, Atiq Ur Rahmn (2019). Analytic Solution to Benjamin-Bona-Mahony Equation by Using Laplace Adomian Decomposition Method. *Matrix Science Mathematic*, 3(1): 01-04.
- Shehata, M.M. (2015) A Study of Some Nonlinear Partial Differential Equations by Using Adomian Decomposition Method and Variational Iteration Method. *American Journal of Computational Mathematics*, 5, 195-203.

- <http://dx.doi.org/10.4236/ajcm.2015.52016>
- Singh, K., Gupta, R.K., Kumar, S. (2011). Benjamin-Bona- Mahony (BBM) equation with variable coefficients similarity reductions and Painleve analysis. *Applied Mathematics and Computation*, 217, 7021-7027
- Talha Achouri, Khaled Omrani, (2009). Numerical solutions for the damped generalized regularized long-wave equation with a variable coefficient by Adomian decomposition method", *Communication in Nonlinear Science and Numerical Simulation* 14:2025– 2033
- Tarig M. Elzaki (2011a). The new integral transform "elzaki transform". *Global Journal of Pure and Applied Mathematics*, 7(1):57–64.
- Tarig M. Elzaki (2011b). Application of new transform "elzaki transform" to partial differential equations. *Global Journal of Pure and Applied Mathematics*, 7(1):65–70.
- Wazwaz A.M. and Gorguis, A. (2004). "An analytic study of Fisher's equation by using Adomian decomposition method," *Applied Mathematics and Computation*, vol. 154, no. 3, pp. 609–620.
- Wazwaz AM, and Mehanna MS. (2010). The combined Laplace-Adomian method for handling singular integral equation of heat transfer. *Inter J of Nonlinear Science*. 10:248-52.
- Wazwaz AM. (2010). the combined Laplace transform-Adomian decomposition method for handling nonlinear Volterra integro-differential equations, *Applied Mathematics and Computation*. 216(4):1304-9.